

# UNCLASSIFIED

AD NUMBER
AD843447
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; 01 JUL 1968. Other requests shall be referred to Commanding Officer, Department of The Army, Fort Detrick, Attn: Technical Release Branch/TIO, Frederick, MD 21701.
AUTHORITY
SMUFD, D/A ltr, 15 Feb 1972

THIS PAGE IS UNCLASSIFIED

AD843447

TRANSLATION NO. 2157

DATE: 19 July 1968

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID, Frederick, Maryland 21701

DEPARTMENT OF THE ARMY  
Fort Detrick  
Frederick, Maryland

DDC

NOV 21 1968

115700

A

x

Delcourt, R. 1967. Démonstration par l'électrophorèse en gel d'acryle de la spécificité du bleu Luxol en tant que réactif des phospholipides (Demonstration by acrylic gel electrophoresis of the specificity of Luxol blue in the reaction with phospholipid). Comptes rendus des séances de la Société de Biologie (Proceedings of the sessions of the Society of Biology) vol. 161, No. 6, pp. 1480-1481.

The phthalocyanides are blue dyes containing a metallic atom in the center of a heterocyclic nucleus. The metallic ion is from the oxide used as a catalyst in the synthesis of the dye (Cram and Hammond (1)). Pearse (2) demonstrated that these dyes bind to phospholipid structures such as mycin and red blood cell stroma. In vitro, these same dyes form insoluble complexes with phospholipids containing choline. Lison (3) reported, without any remarks, a limitation in the specificity of this dye for phospholipids. Salthouse has demonstrated the formation of stoichiometric complexes between Luxol blue and phospholipids containing choline or serine.

We have studied the action of Luxol fast blue on blood proteins and their fractions isolated by acrylic gel electrophoresis using a method we previously described (Delcourt (5)).

1. Action on Fixed Proteins. Several spottings of serum on Whatmann No. 1 paper were stained, after dessication, with an alcoholic solution of Luxol fast blue. The excess stain was eluted from the paper using 5 % acetic acid. In addition, the stain fixed to the serum spots was not soluble in a number of conventional solvents. There was, therefore, co-precipitation.

2. Action on Blood Serum. A saturated solution of stain in cellosolve was prepared. The addition of less than 40  $\mu$ l of this solution to 1 ml of serum stained it without any precipitation occurring. When 40  $\mu$ l or more were used, precipitation occurred. This precipitate was insoluble in water and in the usual solvents. The precipitate was quite certainly the result of the formation of an insoluble complex. There was a negative test for free copper in an aqueous solution of the stain; the addition of sodium hydroxide to the serum mixture did not give the characteristic biuret color.

3. Thin-Layer Chromatography. Thin-layer chromatography did not show any migration or any fractionation.

4. Acrylic Gel Electrophoresis. This confirmed the specific fixation of the lipoproteins: (A) the dye alone, in aqueous solution, migrated rapidly towards the cathode without leaving any material along its path of migration; (B) lipid-free solutions of Cohn fractions migrated without retaining any of the dye; (C) a solution of thromboplastin, probably denatured by phenol but still maintaining intact biological activity, was separated into several bands which stained with Luxol blue and whose lipid nature was confirmed by staining with acetyl Sudan blue; (D) several human and animal sera, pre-stained with Luxol blue, gave from four to five fractions whose color reactions were similar to those for alpha-lipoproteins.

Discussion. → We have confirmed the preferential fixation of Luxol blue to lipoproteins rich in phospholipids, such as alpha-lipoproteins and thromboplastin, and the complete absence of staining with other native or lipid-free proteins. The mode of fixation does not occur at the level of the active groups responsible for electrophoretic mobility of lipoproteins, as

shown by the simultaneous electrophoresis of serum prestained with Sudan blue, Nile blue, and Luxol fast blue. It is not the copper atom that interacts and the method of coupling suggested by Pearse is still possible - the formation of a salt of the phospholipid with the dye.

#### References

- (1) P.J. Cram and G.S. Hammond, *Chimie organique*, Presses Univ. Laval, Quebec, 1963, p. 680.
- (2) A.G. Pearse, *Jour. Path. Bact.*, 1955, 70: 554.
- (3) L. Lison, *Histochimie et cytochimie animales*, Gauthier-Villars, Paris, 1960, p. 524.
- (4) T.N. Salthous, *Nature*, 1962, 195: 187.
- (5) R. Delcourt, *Protides of the Biological Fluids*, 15th Colloquium, Bruges, 1977. To be published.